

**THE STRUCTURE OF NAKED SINGULARITY IN SELF-SIMILAR
GRAVITATIONAL COLLAPSE II**

P. S. Joshi and I. H. Dwivedi*
Tata Institute of Fundamental Research
Homi Bhabha Road, Bombay 400 005
India

*** Permanent Address:**
Institute of Basic Sciences
Agra University
Khandari, Agra, India

Proofs to be sent to:
Dr. Pankaj S. Joshi
Theoretical Astrophysics Group
T.I.F.R. Homi Bhabha Road
Colaba, Bombay 400 005
India

e spacing) 72.27 pt=1 inch

Abstract

Generalizing the results of Joshi and Dwivedi in Commun.Math.Phys. 146, p.333 (1992), it is pointed out that strong curvature naked singularities could occur in the self-similar gravitational collapse of any form of matter satisfying the weak energy condition for the positivity of mass-energy density.

The formation of naked singularities in self-similar gravitational collapse of a perfect fluid with an adiabatic equation of state was analyzed in [1] (to be referred to here as paper I). It was shown that a powerfully strong curvature naked singularity forms in that case from which a non-zero measure set of non-spacelike trajectories come out, which could be locally or globally naked. As there is no proof available as yet for the cosmic censorship hypothesis, which is fundamental to the black hole physics and its applications, such an analysis of collapse scenarios appears inevitable in order to prove, and even to formulate a mathematically rigorous censorship hypothesis. This would allow the features of collapse to be isolated which must be avoided in order to establish cosmic censorship.

Eventhough the form of matter such as a perfect fluid has a wide range of physical applications with the advantage of incorporating the pressure which could be important in the later stages of collapse, it is certainly important to examine if similar conclusions will hold for other reasonable forms of matter. In fact, as pointed out by Eardley [2], the forms of matter such as dust could be an approximation to more fundamental forms, such as a massive scalar field. It is thus conceivable that the naked singularity could be an artifact of the approximation used, rather than being a basic feature of gravitational collapse, and therefore it is important to examine these conclusions for a broader range of matter.

With this purpose, we examine here the final fate of a self-similar collapse to show that conclusions on the occurrence and nature of naked singularity generalize to all forms of matter which satisfy the positivity of energy as characterized by the well-known energy conditions [3]. It follows that the conclusions of paper I are not limited to a specific form of matter such as a perfect fluid.

The non-zero metric components for a spherically symmetric space-time are

$(t, r, \theta, \phi = 0, 1, 2, 3)$

$$g_{00} = -e^{2\nu}, \quad g_{11} = e^{2\psi} \equiv V + X^2 e^{2\nu}, \quad g_{33} = g_{22} \sin^2 \theta = r^2 S^2 \sin^2 \theta \quad (1)$$

where V is defined as above and due to self-similarity ν, ψ, V and S are functions of the similarity parameter $X = t/r$ only. The remaining freedom in the choice of coordinates r and t has been used to set the only off-diagonal term T_{01} of the energy momentum tensor T_{ij} to zero (using comoving coordinates). We assume the matter to satisfy the weak energy condition, i.e.

$$T_{ij} V^i V^j \geq 0 \quad (2)$$

for all non-spacelike vectors V^i . Therefore, $T_0^0 \leq 0$ (i.e $T_{00} \geq 0$), $T_1^1 - T_0^0 \geq 0$, $T_2^2 - T_0^0 \geq 0$. The relevant field equations for a spherically symmetric self-similar collapse of the fluid under consideration are [4],

$$G^0_0 = \frac{-1}{S^2} + \frac{2e^{-2\psi}}{S} (X^2 \ddot{S} - X^2 \dot{S} \dot{\psi} + X S \dot{\psi} + \frac{(S - X \dot{S})^2}{2S}) - \frac{2e^{-2\nu}}{S} (\dot{S} \dot{\psi} + \frac{\dot{S}^2}{2S}) = 8\pi r^2 T^0_0 \quad (3a)$$

$$G^1_1 = \frac{-1}{S^2} - \frac{2e^{-2\nu}}{S} (\ddot{S} - \dot{S} \dot{\nu} + \frac{\dot{S}^2}{2S}) + \frac{2e^{-2\psi}}{S} [-S X \dot{\nu} + X^2 \dot{S} \dot{\nu} + \frac{(S - X \dot{S})^2}{2S}] = 8\pi r^2 T^1_1 \quad (3b)$$

$$G^2_2 = 8\pi T^2_2, \quad G^0_1 = \ddot{S} - \dot{S} \dot{\nu} - \dot{S} \dot{\psi} + \frac{S \dot{\psi}}{X} = T^0_1 = 0 \quad (3c)$$

Using equation (3c), equations (3a) and (3b) can be combined to get

$$\dot{V}(X) = X e^{2\nu} [H - 2] \quad (3d)$$

where $(\cdot)$ is the derivative with respect to the similarity parameter $X = t/r$ and $H = H(X)$ is defined by

$$H = r^2 e^{2\psi} (T^1_1 - T^0_0) \quad (4)$$

For matter satisfying weak energy condition, it follows that $H(X) \geq 0$ for all X . Here we have four field equations with six unknowns. The remaining two equations come from the choice of form of matter one is dealing with. Using (3b) and methods similar to paper I one can see that the singularity at $t = 0, r = 0$ is naked when the equation $V(X) = 0$ has a real simple positive root, i.e.

$$V(X_0) = 0 \tag{5}$$

for some $X = X_0$; (see also Waugh and Lake [6], and the use of such a condition in the numerical simulations by Ori and Piran [6]; and also [5] for the relevance of the same for naked singularity formation in self-similar radiation collapse models); and a non-zero measure of future directed non-spacelike trajectories will escape from the singularity provided

$$0 < H_0 = H(X_0) < \infty \tag{6}$$

These escaping trajectories near the singularity are given by,

$$r = D(X - X_0)^{\frac{2}{H_0 - 2}} \tag{7}$$

Here D is a constant labeling different integral curves, which are the solutions of the geodesic equations, coming out of the naked singularity. It is seen that $H_0 > 0$ will hold if the weak energy condition (2) is satisfied and when the energy density as measured by any timelike observer is positive in the collapsing region near the singularity. In this case, when $H_0 < \infty$, families of future directed non-spacelike geodesics will come out, terminating at the naked singularity in the past. On the other hand, for $H_0 = \infty$, a single non-spacelike trajectory will come out of the naked singularity. This characterizes the formation of naked singularity in self-similar gravitational collapse. Such a singularity will be atleast locally naked and

considerations such as those in paper I can be used to show that it could be globally naked as well provided $V(X_0) = 0$ has more than one real simple positive roots.

We do not go here into a detailed discussion of the sufficient conditions which would ensure the real positive roots for (5) (and hence the existence of naked singularity) in terms of physical parameters involved and the initial data specified prior to the onset of the singularity. However, the existence of several classes of self-similar solutions to Einstein equations, where the gravitational collapse from a regular initial data results into a naked singularity, indicates that such a condition will be realized for a wide variety of self-similar collapse scenarios. For example, the cases of radiation collapse models with a linear mass function [5], and self-similar Tolman-Bondi models [6] are all special cases of the treatment given here. For the case of radiation collapse with the mass function $m(u) = \lambda u$ (where u is the advanced time), the above initial condition corresponds to a restriction on the parameter λ (which is the rate of collapse) given by $0 < \lambda \leq 1/8$. The collapse must result into a naked singularity when λ is in this range, and an event horizon covers the singularity otherwise. Similar restrictions are obtained, ensuring the existence of a real positive root for $V(X) = 0$, for classes of Tolman-Bondi dust models [7] and adiabatic perfect fluid [1] (in which case this turns out to be a forth order algebraic equation). In general, for a given form of matter, the existence of a real positive root of (5) would put a restriction on the range of values of physical parameters involved. One could also treat condition (5) for the existence of a naked singularity as an initial value problem for the differential equation (3d). That is, for a given form of matter one could solve this first order differential equation governing $V(X)$, with the initial value $V(X_0) = 0$, for some real positive value $X = X_0$. It is possible that the regular initial data specified for a realistic gravitational collapse might just always avoid this initial condition and could result into a black hole. On the

other hand, all the initial data sets satisfying this condition will necessarily evolve into a naked singularity regardless of the form of matter involved. It is thus seen that a naked singularity would form in the collapse of a wide range of matter forms satisfying the weak energy condition.

Apart from non-zero measure of non-spacelike families coming out, the curvature strength of naked singularity provides an important test of its physical significance. This was calculated in paper I along all the non-spacelike geodesics terminating at the naked singularity in the past to show that this is a strong curvature singularity in a powerful sense. Even for the case of the general form of matter considered here, this turns out to be a strong curvature naked singularity. We consider here only the radial null geodesics coming out which are given by,

$$\frac{dt}{dr} = e^{\psi-\nu} \quad (8)$$

The curvature strength of the singularity is measured by evaluating the limit of $k^2 R_{ij} V^i V^j$ along these trajectories near the singularity, which is given in the present case by,

$$\lim_{k \rightarrow 0} k^2 R_{ij} V^i V^j = \frac{4H_0}{(2 + H_0)^2} > 0 \quad (9)$$

It follows that this is a strong curvature naked singularity in the sense that the volume forms defined by all possible Jacobi vector fields vanish in the limit of approach to the naked singularity, in which case the space-time may not admit any continuous extension through the singularity [8].

REFERENCES

1. P.S.Joshi and I.H.Dwivedi, Commun. Math. Phys. 146,333(1992).
2. D. M. Eardley in "Gravitation in Astrophysics", Plenum Publishing Corporation Edited by B. Carter and J. B. Hartle 223 (1987).
3. S.W.Hawking and G.F.R.Ellis, "The Large Scale Structure of Space-time", CUP, Cambridge (1973).
4. M.E. Cahill and A.HY.Taub, Commun.Math.Phys. 21, 1(1971).
5. I.H. Dwivedi and P.S. Joshi, Class. Quantum Grav. 6, 1599 (1989); 8, 1339 (1991).
- 6 B.Waugh and K.Lake, Phys. Rev. D38, 4, 1315 (1988); A.Ori and T.Piran, Phys. Rev. D42, 4, 1068 (1990).
- 7 I.H.Dwivedi and P.S.Joshi, Class. Quantum Grav. 9, L69 (1992).
8. F.J.Tipler, C.J.S.Clarke and G.F.R.Ellis, "General Relativity and Gravitation", Vol 2, (ed.A.Held), New York: Plenum Press, 97 (1980).